

The Endothelium in Clinical Practice

Source and Target of Novel Therapies

edited by

Gabor M. Rubanyi


*Berlex Biosciences, Inc.
Richmond, California*

Victor J. Dzau

*Brigham and Women's Hospital
and Harvard Medical School
Boston, Massachusetts*

MARCEL DEKKER, INC.

NEW YORK • BASEL • HONG KONG



RC 691
- 4
- E54
1997

ISBN: 0-8247-9809-0

The publisher offers discounts on this book when ordered in bulk quantities. For more information, write to Special Sales/Professional Marketing at the address below.

This book is printed on acid-free paper.

96-49906

Copyright © 1997 by MARCEL DEKKER, INC. All Rights Reserved.

Neither this book nor any part may be reproduced or transmitted in any form or by any means, electronic or mechanical, including photocopying, microfilming, and recording, or by any information storage and retrieval system, without permission in writing from the publisher.

MARCEL DEKKER, INC.
270 Madison Avenue, New York, New York 10016

Current printing (last digit):
10 9 8 7 6 5 4 3 2 1

PRINTED IN THE UNITED STATES OF AMERICA

Mar
faci
and
had
per

No
key
Th
suc
ce
pl
"t
cc

ye
B
te

Ryan

soluble com-
pl Int 1994;

udolph, AR,
ibition using
ion f pig-to-
Xenotrans-

I, Levin, JL,
racute rejec-

ient and pig-
Congress for

The Receptor for Advanced Glycation Endproducts: Implications for the Development of Diabetic Vascular Disease

Osamu Hori, Shi Du Yan, and Ann Marie Schmidt

Columbia University College of Physicians and Surgeons, New York, New York

I. INTRODUCTION

Proteins or lipids exposed to aldose sugars undergo nonenzymatic glycation and oxidation (1-7). The ultimate products of these interactions are called the advanced glycation endproducts (AGEs). Earlier, however, reversible adducts known as Schiff bases/Amadori products form in the course of exposure to aldoses. These products, distinct from AGEs, are reversibly formed and can dissociate to the native proteins on restoration of normoglycemia (4). Clinically, the reversible products have been used to assess the adequacy of glycemic control. The best known of these moieties used in this setting is hemoglobin A_{1c} (8). In contrast, AGEs are irreversibly formed. Although a heterogeneous group of compounds, they nevertheless share in common certain characteristics such as yellow-brown color, the tendency to form cross links, the ability to generate reactive oxygen intermediates, fluorescence, and the ability to recognize specific receptors on cellular surfaces (9-13). AGEs accumulate in the plasma and vessel wall during normal aging, but they form to an accelerated degree in patients with diabetes.

A number of groups have developed specific reagents (both polyclonal and monoclonal antibodies) to detect the presence of AGEs in vivo (14-17). For example, we have identified increased immunoreactivity for AGEs in the vasculature of diabetic patients versus that in age-matched,

nondiabetic control individuals (18). Similarly, others have found increased immunoreactivity for AGEs in the vessel wall and skin of diabetic patients (19–22). The results of these studies are consistent with the hypothesis that formation of AGEs correlates with the development of diabetic complications. As well, previous studies have suggested a possible link between the accumulation of AGEs in the vasculature and the development of vascular disease which occurs to an accelerated extent in diabetic individuals. For example, Nakamura and colleagues demonstrated increased immunoreactivity for AGEs in coronary atheromas of patients with diabetes (22).

It should be noted that in addition to normal aging (1,2,4) and diabetes (1–8), the development of proteins/lipids modified by irreversible nonenzymatic glycation may occur under other conditions which favor their generation. For example, AGEs may form on long-lived proteins, particularly in the setting of renal failure. In this context, under physiological conditions, the plasma protein β_2 -microglobulin is readily cleared from the circulation. However, in the presence of its markedly delayed clearance characteristic of renal failure, β_2 -microglobulin may undergo AGE formation (23), presumably as a consequence of its prolonged half-life in the setting of renal failure. In marked contrast to the native form, AGE- β_2 -microglobulin is postulated to be important in the pathogenesis of the inflammatory disorder dialysis-related amyloidosis (DRA) that is associated with chronic renal failure and dialysis (24).

Another condition in which amyloid deposition may predispose to the development of AGE formation is in Alzheimer's disease (25). In this disorder, AGE formation may form on the components of intracellular neurofibrillary tangles (26–28) as well as in extracellular amyloid- β -peptide accumulations (29). The presence of these AGE-modified moieties has been postulated to be an important component of enhanced oxidant stress, which is likely involved, at least in part, in the pathogenesis of this disorder (26,28).

Taken together, these data suggest that AGEs may form under conditions of hyperglycemia, normal aging, delayed protein turnover (such as in renal failure), or extensive protein accumulation (such as in amyloidoses). In this chapter, we detail some of the biological properties of these structures with respect to their likely contributory role in the development of vascular/cellular pathology.

II. IDENTIFICATION OF CELLULAR RECEPTORS FOR AGEs

Earlier studies identified the ability of AGE-modified ligands to bind to cellular surfaces (4,30) in a specific and saturable manner. Although a number of cellular binding sites for AGEs have been identified (11,31,32), the

The Receptor for Ad

best characterized of products (RAGE) (1) fast-pressure liquid ionization media, RAGE bovine lung. At 1 lactoferrin-like poly confined to consid

Molecular cloning of a member of the The predicted hydrophobic suggests that RAGE one "V"-type immun This is followed by highly charged cy signal sequence of human counterpar hypothesis that RAGE binding of 125 I-AGE RAGE but not to RAGE-transfected min to immobilize (11,13,16) with K RAGE IgG (11,1

Given the fact family, we postul




Figure 1 Hydrophobicity plot generated by the Woods program. NH₂, amino terminal sites; TM, transmembrane; Ig, immunoglobulin.

increased
ic patients
thesis that
complica-
between the
of vascular
duals. For
immuno-
etes (22).
d diabetes
nonenzym-
air genera-
icularly in
onditions,
irculation.
racteristic
(23), pre-
ig of renal
globulin is
tory disor-
onic renal

ose to the
this disor-
tracellular
- β -peptide
s has been
ess, which
er (26,28).
der condi-
(such as in
yloidoses).
ese struc-
opment of

to bind to
igh a num-
1,32), the

best characterized of these to date is the receptor for advanced glycation endproducts (RAGE) (11). By sequential chromatography on hydroxylapatite, fast-pressure liquid chromatography (FPLC) Mono S, and FPLC gel filtration media, RAGE was purified to homogeneity from detergent extracts of bovine lung. At the same time, another AGE binding polypeptide, lactoferrin-like polypeptide (LF-L) was also described (11). This chapter is confined to consideration of RAGE.

Molecular cloning studies (13) have revealed that RAGE is a newly identified member of the immunoglobulin superfamily of cell surface molecules. The predicted hydropathy plot of the bovine cDNA for RAGE (Fig. 1) suggests that RAGE contains a 332-amino acid extracellular domain with one "V"-type immunoglobulin domain followed by two "C"-type domains. This is followed by a hydrophobic transmembrane spanning domain and a highly charged cytosolic tail. Preceding the above domains is a putative signal sequence of 22 amino acids. Of note, the bovine, murine, rat, and human counterparts of RAGE are remarkably similar. Critical to the hypothesis that RAGE was an AGE binding protein was the demonstration of binding of 125 I-AGE albumin to 293 cells transfected with the cDNA for RAGE but not to cells transfected with irrelevant cDNA (13). Binding to RAGE-transfected 293 cells was similar to that of radiolabeled AGE albumin to immobilized RAGE, endothelial cells, or mononuclear phagocytes (11,13,16) with $K_d \approx 50$ –100 nM and inhibitable in the presence of anti-RAGE IgG (11,13,16).

Given the fact that RAGE was a member of the immunoglobulin superfamily, we postulated that under physiological conditions, RAGE might




Figure 1 Hydropathy plot of bovine RAGE. As generated by the Hopp and Woods program of intelligenetics, this hydropathy plot of RAGE was generated. NH₂, amino terminus; COOH, carboxyl-terminus; CHO, N-linked glycosylation sites; TM, transmembrane domain; S-S, disulfide-linked cysteine residues; IgV and IgC, immunoglobulin-like variable and constant domains, respectively.

function as a cell-cell adhesion molecule or receptor for cytokine(s) or growth factor(s). In this context, we have demonstrated that under the specific condition of perinatal development, RAGE is highly expressed in the developing central nervous system of the rat. Its specific interaction with the polypeptide amphoterin appears to mediate neurite outgrowth in *in vitro* assays designed to study this phenomenon (33). Another recently identified putative ligand of RAGE, a molecule with $M_r \approx 12$ kD, is a newly identified member of a family of proinflammatory cytokines (33). Detailed analysis of the consequences of its interaction with RAGE are in progress.

In addition to the presence of RAGE on endothelial cells and mononuclear phagocytes, a survey of normal tissue has identified its presence on other cell types such as mesangial cells and certain neurons (34). In addition, RAGE is also present in the vascular smooth muscle (34). As stated above, RAGE is also present to a significant degree in the developing nervous system in embryonic and postnatal (through P17) cortical neurons (33). These data lend further support to the hypothesis that AGEs may in fact be pathological ligands of this receptor, which under homeostatic conditions, act to mediate such functions as normal development and, potentially, other not yet identified phenomena.

In this chapter, we concentrate on the implications of AGE-RAGE interaction on mononuclear phagocytes and endothelial cells.

III. INTERACTION OF AGEs WITH MONONUCLEAR PHAGOCYTE RAGE: INITIATION OF MONONUCLEAR PHAGOCYTE MIGRATION AND ACTIVATION

Previous studies have identified the presence of RAGE on mononuclear phagocytes by immunohistochemistry (16) and by *in situ* hybridization (S.D. Yan and A.M. Schmidt, unpublished observation). Radioligand binding studies with 125 I-AGE albumin demonstrated specific saturable binding, with $K_D \approx 80$ nM, comparable to that observed on immobilized RAGE (11). Consistent with the concept that RAGE was important in mediating the interaction of AGEs with mononuclear phagocytes (MPs), binding was largely blocked in the presence of anti-RAGE IgG.

We hypothesized that soluble AGEs, such as those found circulating on modified plasma proteins or on the surface of diabetic red blood cells (35), for example, might initiate monocyte migration. In this context, when soluble AGEs (either AGE albumin or AGEs immunisolated from diabetic plasma using affinity-purified anti-AGE IgG) were placed in the lower compartment of microchemotaxis chambers, monocyte migration was promoted in a dose-dependent manner (16). In contrast, native albumin was without effect. Furthermore, that this was true chemotaxis was demonstrated by

further studies with AGE- β_2 microglobulin. Migration of monocytes in the presence of AGEs was blocked by the addition of soluble RAGE (16). In contrast, binding of AGEs to RAGE was initiated by the

In addition to the role of RAGE in the development of the AGI, the role in the development of the AGI is significant, they study this hypothesis on the upper surface of monocyte migration in the lower compartment of albumin, monocyte migration increased, as expected the upper surface compartment was restored in monocyte migration studies preincubated with FMLP was restored to the upper surface. Similarly, when monocyte migration was restored (16).

To further test the hypothesis, we utilized RAGE, assays directly to native albumin coated with native albumin migration was (left panel). In coated with AGEs, short paths of

Analogous to this concept. P either native or implanted subcutaneous tissue of

line(s) or under the pressed in interaction growth in recently s a newly Detailed progress. id mono- sence on . In addi- As stated eveloping l neurons s may in tic condi- d, poten-

E-RAGE

LEAR

ionuclear ridization and bind- binding, GE (11). ating the ding was

dating on xells (35), when sol- i diabetic wer com- promoted s without trated by

further studies with checkerboard analysis. Similar results were observed with AGE- β_2 microglobulin with respect to true monocyte chemotaxis (24). Migration of monocytes in response to soluble AGE albumin was inhibited in the presence of anti-RAGE IgG (or F(ab')₂ fragments) or in the presence of soluble RAGE (sRAGE, the extracellular two thirds of the molecule [16]). In contrast, blockade of RAGE was without effect on monocyte migration initiated by the chemotactic peptide F-met-leu-phe (FMLP).

In addition to soluble AGEs, AGEs may also form on the long-lived proteins of the subendothelium, an area postulated to have an important role in the development of vascular lesions, such as atherosclerosis (36). In order for the AGEs formed/deposited in this area to be of pathological significance, they would likely need to attenuate monocyte migration. To study this hypothesis, we immobilized either AGE albumin or native albumin on the upper membranes of microchemotaxis chambers and tested monocyte migration in response to the chemotactic peptide FMLP placed in the lower compartment. When the upper surface was coated with native albumin, monocyte migration in response to increasing doses of FMLP was increased, as expected. However, when AGE albumin was immobilized on the upper surface, monocyte migration in response to FMLP in the lower compartment was attenuated in a dose-dependent manner (16). The central role of monocyte RAGE in mediating these interactions was demonstrated in studies employing RAGE blockade. When monocytes were preincubated with anti-RAGE F(ab')₂, monocyte migration in response to FMLP was restored despite the presence of immobilized AGE on the upper surface. In contrast, nonimmune fragments were without effect. Similarly, when the AGE-coated surface was pretreated with sRAGE, monocyte migration in response to FMLP in the lower compartment was restored (16).

To further test the hypothesis that when migrating MPs encounter immobilized RAGE, their migration is slowed, we performed phagokinetic track assays directly to visualize monocyte migration on either AGE albumin or native albumin-coated surfaces. When MPs were placed on coverslips coated with native albumin that had been overlaid by colloidal gold, their migration was evident with long paths of migration demonstrated (Fig. 2, left panel). In marked contrast, when MPs were placed on coverslips coated with AGE albumin, their migration was slowed, as evidenced by short paths of migration (Fig. 2, right panel).

Analogous experiments were performed *in vivo* to further demonstrate this concept. Polytetrafluoroethylene (PTFE) tubes were impregnated with either native rat serum albumin or AGE rat serum albumin and then implanted subcutaneously onto the backs of rats. When removed after 4 days, the tissue of rats treated with PTFE tubes to which had been absorbed




Figure 2 Phagokinetic track assay. MPs were allowed to migrate for 6 hr on matrices of native albumin (left panel) or AGE albumin (right panel) which had been coated with colloidal gold particles. The dark tracks made by locomoting MPs are visualized by darkfield microscopy. $\times 125$

native albumin revealed minimal inflammatory infiltrate except at the tissue-tube interface (Fig. 3, left panel). In contrast, however, tissue from rats treated with AGE-impregnated PTFE tubes revealed a considerable inflammatory infiltrate, with mononuclear cells observed moving through the interstices of the tubes (Fig. 3, right panel). Many of the monocytes appear to have undergone activation, as evidenced by leukocytoclastic changes.

These data are consistent with the concept that AGEs may promote monocyte activation. In fact, previous studies have shown that AGE-monocyte interaction results in their activation (37–38), as manifested by production of cytokines such as tumor necrosis factor- α (TNF- α) and interleukin-1 (IL-1) as well as the growth factor platelet-derived growth factor (PDGF).

Taken together, these data suggest a model in which soluble AGEs, such as those circulating in modified plasma proteins or those on the surface of diabetic red blood cells, may attract monocytes and mediate their migra-



6 hr on
hich had
ing MPs

at the
ue from
derable
through
nocytes
oclastic

romote
AGE-
sted by
(α) and
growth

Es, such
rface of
migma-



Figure 3 Implantation of PTFE tubes with absorbed AGE albumin into rats. PTFE tubes were impregnated with either native albumin (left panel) or AGE albumin (right panel), implanted subcutaneously into the backs of rats, and removed after 4 days and stained with hematoxylin and eosin. $\times 110$.

tion. In contrast, immobilized AGEs, such as those formed/deposited in the vascular wall, may retard the migration of monocytes, a process which may ultimately lead to their activation, a process which may predispose to the development of vascular lesions.

IV. INTERACTION OF AGEs WITH ENDOTHELIAL CELL RAGE: CLEARANCE OF AGEs BY ENDOTHELIAL CELLS, INDUCTION OF CELLULAR OXIDANT STRESS, AND ENHANCED EXPRESSION OF VASCULAR CELL ADHESION MOLECULE-1

A. Endothelial Cell Handling of AGEs: Clearance Studies and Experiments Assessing Cellular Activation

Previous studies have identified the presence of RAGE in the vascular endothelium by immunohistochemistry using anti-RAGE IgG and by *in situ* hybridization (34). The endothelium is in a critical position to mediate the effects of circulating AGEs, either those that form on circulating plasma proteins or those that deposit/form on the surface of diabetic red blood cells with the rest of the cell. In addition to its potential role in processing/removing AGEs from the intravascular space, the endothelium may also function to mediate certain perturbations that are a consequence of their interaction with AGEs, all of which may alter the cellular phenotype, thereby predisposing to vascular dysfunction. In this context, our earliest studies on AGE-endothelial interaction demonstrated that when cultured endothelial cells (ECs) were exposed to AGEs, critical cellular properties were altered resulting in a procoagulant and more permeable phenotype (30). Specifically, we observed suppression of thrombomodulin, sustained induction of low levels of tissue factor, and increased diffusional transit of macromolecular solutes across the EC monolayer (30). The central role of RAGE in mediating the interaction of AGEs with the endothelium is suggested by studies which demonstrated blockade of binding of radiolabeled AGEs to cultured ECs (11) by anti-RAGE IgG(11).

It was therefore most important in this context to study the effects of AGEs infused into animals. Extensive studies with ¹²⁵I-AGE albumin administered intravenously into mice revealed that in contrast to radiolabeled native albumin, radiolabeled AGE albumin demonstrated an initial rapid phase of clearance, with greater than 50% of the protein being cleared from the vascular space within minutes after administration (39). Studies assessing tracer accumulation in the organs demonstrated that AGEs accumulated in the most vascular organs (39). Studies utilizing pretreatment of the animals with either anti-RAGE IgG or sRAGE resulted in a marked

The Receptor for

delay in AGE c
interfering with

Electronmicro
gold revealed th
interact with th
the EC layer,
abluminal surfa
this process was
whereas nonim

The importa
nonradioactive
leukin-6, again
(39). Taken to
interacting with
tion of cytokin
lesions. The ce
strated by stud
to cell surface
intravascular
RAGE utilizir
case, induction

B. Induction AGEs vi

During the pr
tive oxygen in
at least in part
tion might al
stress. Initial
min to cultur
reactive subst
activation of
min was with
dant stress, w
rable results
cells isolated

To better
AGEs delive
into normal
for heme o
phoretic mc

sited in
is which
pose to

LL

LESS,
JL

id

vascular
d by in
mediate
ulating
etic red
role in
thelium
quence
pheno-
xt, our
it when
cellular
meable
odulin,
fusional
he cen-
e endo-
ding of

fects of
nin ad-
labeled
ul rapid
cleared
Studies
is accu-
nent of
marked

delay in AGE clearance, suggesting that these reagents were capable of interfering with the interaction of circulating AGEs with EC RAGE (39).

Electronmicroscopic studies utilizing AGE albumin linked to colloidal gold revealed that in a murine coronary vasculature model, AGEs quickly interact with the EC surface, are subsequently endocytosed, and traverse the EC layer, in part by transcytosis, ultimately being released at the abluminal surface (39). As with the radiolabeled AGE clearance studies, this process was significantly inhibited in the presence of anti-RAGE IgG, whereas nonimmune IgG was without effect.

The importance of these results is stressed by studies in which infused, nonradioactive AGEs into mice resulted in increased mRNA for interleukin-6, again a process inhibited in the presence of RAGE blockade (39). Taken together, these studies suggest that AGEs are capable of interacting with EC RAGE, with potential consequences, including activation of cytokines, which may predispose to the development of vascular lesions. The central role of RAGE in mediating these effects is demonstrated by studies in which blockade of RAGE—either by blocking access to cell surface RAGE with anti-RAGE IgG or by binding to AGEs in the intravascular space thereby limiting their interaction with cell surface RAGE utilizing sRAGE—inhibits not only AGE clearance but, in this case, induction of, for example, mRNA for IL-6 transcripts.

B. Induction of Oxidant Stress in the Vasculature by Exposure to AGEs via a RAGE-Dependent Pathway

During the process of AGE formation, oxidation and generation of reactive oxygen intermediates are formed (9,10). We therefore postulated that, at least in part, one of the main mechanisms by which AGE-RAGE interaction might alter/perturb cellular properties is via generation of oxidant stress. Initial studies in tissue culture revealed that exposure of AGE albumin to cultured ECs resulted in the production of thiobarbituric acid-reactive substances (TBARS), induction of heme oxygenase mRNA, and activation of the transcription factor NF- κ B (17). In contrast, native albumin was without effect. All of these markers, indicative of enhanced oxidant stress, were inhibited in the presence of anti-RAGE IgG (17). Comparable results were obtained when diabetic red blood cells, not red blood cells isolated from normal individuals, were exposed to cultured ECs (35).

To better test the hypothesis that ligation of cell surface RAGE by AGEs delivered oxidant stress to the tissues, AGE albumin was infused into normal mice resulting in increased tissue TBARS, increased mRNA for heme oxygenase, and activation of NF- κ B as assessed by electrophoretic mobility shift analysis of liver nuclear extracts (17,40). Again,

native albumin was without significant effect, and these processes in AGE-treated animals were inhibited by preinfusion of anti-RAGE IgG.

To detect these malondialdehyde-reactive epitopes in the vasculature itself, AGE albumin or native albumin was administered intravenously into rats, and after 1 h, samples were taken for immunohistochemical analysis using antibody that specifically recognizes malondialdehyde epitopes (17). Analysis of the lungs revealed vascular staining (in the ECs and vascular smooth muscle) of AGE-treated animals (Fig. 4, left panel) suggestive of enhanced oxidant stress. In contrast, no staining was observed in the pulmonary vasculature of animals treated with native albumin (Fig. 4, right panel). In the former case, nonimmune IgG was similarly without effect (17).

These studies were important to identify the site(s) where AGEs deliver oxidant stress after their infusion. Taken together, the data suggest that



Figure 4 Immunohistological detection of malondialdehyde-reactive epitopes in the vasculature of rats infused with AGE versus native rat serum albumin. Rats were infused with either AGE rat serum albumin (left panel) or native rat serum albumin (right panel). After 60 min, the animals were killed and lungs processed for the detection of malondialdehyde using indirect immunoalkaline phosphatase with antibody generously supplied by Dr. Joseph Witztum, University of California at San Diego. $\times 400$.

The Receptor for

AGEs interact with this interaction: multiple potent consequences

The implications reaching in this linked to oxidant stress probably involve such entities as which may be in context, we consider the role of the RAGE molecule with EC RAGE.

C. AGE-RAGE

It is likely that the course of vascular disease in controls. On the enhanced endothelium, which is part of the vessel wall. In part, the transmembrane sensitive pathway with their structural and functional

In the setting of atherosclerosis, the expression of VCAM-1 has been shown to be increased in diabetic with

Our data is consistent with the findings from cells, ELISA, and Western blot analysis. VCAM-1 compared with control to assess the function. We treated ECs with AGEs and observed increased binding of AGEs to ECs. This binding bears the consequences of increased oxidant stress and increased expression of RAGE, which is inhibited in the



AGEs interact with EC surface RAGE, and one of the consequences of this interaction is enhanced oxidant stress to the tissues, a process with multiple potential consequences. In this setting, one of the most important consequences may be activation of NF κ B.

The implications of activation of the transcription factor NF κ B are far reaching in this setting. Activation of this transcription factor has been linked to oxidant-sensitive mechanisms (40). Furthermore, its activation is probably important in the ultimate regulation of increased transcription of such entities as adhesion molecules, cytokines, or growth factors; all of which may be important in the pathogenesis of vascular lesions (41). In this context, we considered that enhanced expression of EC vascular cell adhesion molecule-1 (VCAM-1) might be a result of the interaction of AGEs with EC RAGE.

C. AGE-RAGE Interaction Increases EC Expression of VCAM-1

It is likely that many factors contribute to the increased and accelerated course of vascular disease in diabetic individuals compared with normal controls. One of the critical steps in this process, however, is likely to be the enhanced expression of adhesion molecules on the vascular endothelium, which results in the adherence of circulating blood cells to the vessel wall. In this context, previous studies have indicated that, at least in part, the transcriptional regulation of VCAM-1 is controlled by oxidant-sensitive pathways (42). We, therefore, considered whether AGEs interacting with their cellular receptor RAGE might result in increased expression and functional activity of VCAM-1.

In the setting of diabetes, VCAM-1 is of particular interest, as its expression has been identified in the early phase of experimental hypercholesterolemia-induced atherosclerosis (43-44). Furthermore, enhanced VCAM-1 expression has been shown in the vessel wall of rabbits rendered diabetic with alloxan (45), as well as in human atherosclerotic lesions (46).

Our data indicated that when either AGE albumin or AGEs immunoisolated from diabetic plasma were exposed to cultured human endothelial cells, ELISA of cellular extracts revealed increased cell-associated VCAM-1 compared with ECs treated with nonglycated proteins (47). In order to assess the functional implications of increased VCAM-1 on the cell surface, we treated ECs with either AGE or native albumin and subsequently performed binding assays using ⁵¹chromium-labeled Molt-4 cells (cells which bear the counterligand for VCAM-1, VLA-4) and demonstrated significantly increased binding in the presence of AGEs (47). In both cases, the increased expression and function of cell-associated VCAM-1 was largely inhibited in the presence of RAGE blockade but not by control interven-

tions (47). In addition, both processes were inhibited in the presence of antioxidants such as N-acetyl cysteine. Consistent with these data, Northern analysis and nuclear run-on analysis demonstrated increased transcription of VCAM-1 in the presence of AGE albumin, not native albumin, a process which is inhibited by anti-RAGE IgG.

Since previous studies have identified binding sites for NF κ B in the promoter of VCAM-1, which have been shown to participate in the regulation of VCAM-1 in response to such factors as cytokines (48–49), at least in part, by oxidant-sensitive mechanisms. As demonstrated by electrophoretic mobility shift assay, in comparison with native albumin, nuclear extracts prepared from ECs exposed to AGE albumin resulted in an increased DNA binding activity for NF κ B site in the VCAM-1 promoter (Fig. 5, lanes 2 and 3, 6, and 10, respectively). This binding was blocked in a dose-dependent manner in the presence of anti-RAGE IgG (Fig. 5, lanes 11 and 12) but not in the presence of nonimmune IgG (Fig. 5, lane 13). Similarly, pretreatment with N-acetylcysteine resulted in inhibition of AGE-induced activation of NF κ B (Fig. 5, lane 14). Supershift studies using antibodies to p50 and p65 demonstrated that both members of the NF κ B are involved in binding to specific sites in the VCAM-1 promoter (Fig. 5, lanes 7–9). These data suggest that at least of the mechanisms by which AGEs induce enhanced expression of VCAM-1 is via activation of this transcription factor.

Consistent with these *in vitro* studies, infusion of AGE albumin into normal mice resulted in increased immunoreactivity for VCAM-1 in the lung vasculature (Fig. 6, right panel). In contrast, infusion of native albumin was without effect (Fig. 6, left panel).

In addition to increased expression of cell-associated VCAM-1 in the presence of AGEs, one of the further interesting findings in this setting was the detection of VCAM-1 antigen in the supernatants of AGE-treated (as well as TNF- α -treated) ECs (47). We therefore inquired whether patients with diabetes had increased levels of soluble VCAM-1 (sVCAM-1) in their plasma compared with age-matched controls. In both cases, only patients with renal insufficiency were excluded from analysis. The level of sVCAM-1 in the plasma of diabetic patients was found to be 1,115 ng/ml compared with 632 ng/ml in the plasma of nondiabetic controls ($P < .05$). These data paralleled the changes in the levels of AGEs, as measured by AGE albumin equivalent in diabetic plasma compared with control plasma (244 ng/ml versus 157 ng/ml AGE albumin equivalents, $P < .05$) using an ELISA utilizing affinity-purified anti-AGE IgG (16,17,47).

Further analysis of sVCAM-1 in both AGE-treated EC supernatants and patient plasma demonstrated in immunoblotting studies that the immunoreactive material for VCAM-1 is intact, with an $M_r \approx 110$ kD. Fur-

1 2 3



TNF- α
Native Albumin
AGE Albumin
100x

Figure 5 B promoter of VCAM-1 contains NF κ B binding sites. Lane 2, nuclear extracts of ECs treated with AGE albumin; lane 3, ECs were p

presence of
a, North-
transcrip-
tumin, a

B in the
e regula-
it least in
electro-
, nuclear
in an in-
promoter
ocked in
. 5, lanes
lane 13).
ition of
ies using
ie NF κ B
(Fig. 5,
y which
1 of this

min into
1 in the
ve albu-

1 in the
ting was
ated (as
patients
in their
patients
VCAM-
mpared
ese data
iE albu-
14 ng/ml
ELISA

matants
hat the
D. Fur-




Figure 5 Binding of endothelial cell nuclear proteins to NF- κ B binding sites in the promoter of VCAM-1: analysis by electrophoretic mobility shift assay. All lanes contain labeled probe as follows: lane 1, ECs were incubated with TNF- α for 6 hr; lane 2, nonglycated albumin; lanes 3, 6, and 10, AGE albumin; lanes 7-9, nuclear extracts of AGE-treated cells with anti-p50 IgG, anti-p65 IgG, or both anti-p50 IgG and anti-p65 IgG together, respectively. In lanes 11 and 12, ECs were pretreated with anti-RAGE IgG (70 vs 7 μ g/ml, respectively) prior to the addition of AGE. In lane 13, ECs were pretreated with nonimmune IgG prior to AGE, and in lane 14, ECs were pretreated with N-acetylcysteine prior to AGE albumin.




Figure 6 Infusion of AGE albumin induces vascular expression of VCAM-1 in the pulmonary vasculature of mice. Mice were infused with either native albumin (left panel) or AGE mouse serum albumin (right panel). After 30 min, the animals were killed, lungs harvested, and immunostaining with antimurine VCAM-1 antigen performed. $\times 340$.

thermore, no degradation was observed on the immunoblots (50), suggesting that VCAM-1 was released/cleaved from the cell surface in an intact form.

We hypothesized that elevation of sVCAM-1 might be, at least, a marker of endothelial perturbation in diabetic patients compared with control individuals and that its measurement might in fact be a marker of the effectiveness of therapeutic intervention. To further test this concept, we found that the levels of sVCAM-1 in diabetic patients with microalbuminuria were elevated compared with levels of sVCAM-1 in normoalbuminuric diabetic individuals (757 ng/ml vs 505 ng/ml, $P < 0.05$ [50]). The presence of microalbuminuria is of particular importance, since its presence in the diabetic patient has been linked to an increased incidence of cardiovascular morbidity and mortality, and it is considered a marker of diffuse vascular hyperpermeability (51–52). In fact, in our study, 5 of 12 patients with microalbuminuria had evidence of cardiovascular disease, whereas none of 8 patients with normoalbuminuria demonstrated abnor-

The Receptor for

malities of the ca data suggest that lature may be, a interacting with c action may be the wall, which is a p lar disease.

V. BLOCKAI INTERVE COMPLIC

These data sugg receptor RAGE environment fav demonstrated t ment of monoc immobilized A setting. RAGE of AGEs from AGEs results vasculature. On likely to be the data further su for therapeutic Indeed, recent angiogenic pro vascular milieu Taken togethe interaction rep betic vascular

In this cont sRAGE into c (TBIR) obser dysfunction (5

In conclus RAGE intera and accelerate role of RAGE dissecting the or its other li

malities of the cardiovascular system, $P < .05$ (50). Taken together, these data suggest that the enhanced presence of VCAM-1 in diabetic vasculature may be, at least in part, a reflection of the interaction of AGEs interacting with cell surface RAGE. One of the consequences of this interaction may be the increased adherence of mononuclear cells to the vascular wall, which is a process likely pivotal in the development of diabetic vascular disease.

V. BLOCKADE OF RAGE AS A POTENTIAL TARGET FOR INTERVENTION IN THE DEVELOPMENT OF VASCULAR COMPLICATIONS IN DIABETES

These data suggest that the interaction of AGEs with their cell surface receptor RAGE may be of central importance in the creation of an environment favorable to the development of vascular lesions. We have demonstrated that AGE-RAGE interaction is involved in the development of monocyte migration, as well as monocyte retention at sites of immobilized AGEs. Activation of monocytes is a likely sequela in this setting. RAGE has been demonstrated to be important in the clearance of AGEs from the intravascular space, and engagement of RAGE by AGEs results in the generation of enhanced oxidant stress in the vasculature. One of the consequences of this enhanced oxidant stress is likely to be the enhanced expression of cell-associated VCAM-1. These data further suggest that blockade of RAGE might ultimately be a target for therapeutic intervention in the setting of diabetic vascular disease. Indeed, recent findings have suggested that sVCAM-1 may have angiogenic properties, a situation to be of likely detriment in the diabetic vascular milieu, especially, for example, in such sites as the retina (53). Taken together, these findings suggest that blockade of AGE-RAGE interaction represents a logical target for therapeutic intervention in diabetic vascular disease.

In this context, our recent studies have demonstrated that infusion of sRAGE into diabetic rats reversed the increased tissue-blood isotope ratio (TBIR) observed in these animals, which is a marker of endothelial cell dysfunction (54-56).

In conclusion, these data suggest that further assessment of AGE-RAGE interaction and RAGE blockade in such models as wound healing and accelerated atherosclerosis will likely provide insight not only into the role of RAGE as a potential therapeutic target but will also aid in further dissecting the mechanisms set into motion on ligation of RAGE by AGEs or its other ligands.

l in the
in (left
ls were
antigen

ggest-
intact

ast, a
h con-
of the
ot, we
calbu-
calbu-
) The
pres-
nce of
ker of
of 12
sease,
ibnor-

REFERENCES

1. Ruderman N, Williamson J, Brownlee M. Glucose and diabetic vascular disease. *FASEB J* 1992; 6:2905-2914.
2. Baynes J. Role of oxidative stress in development of complications in diabetes. *Diabetes* 1991; 40:405-412.
3. Sell D, Monnier V. Structure elucidation of senescence cross-link from human extracellular matrix: implication of pentoses in the aging process. *J Biol Chem* 1989; 264:21597-21602.
4. Brownlee M, Cerami A, Vlassara H. Advanced glycosylation endproducts in tissue and the biochemical basis of diabetic complications. *N. Engl. J. Med.* 1988; 318:1315-1320.
5. Bucala R, Makita Z, Koschinsky T, Cerami A, Vlassara H. Lipid advanced glycosylation: pathway for lipid oxidation in vivo. *Proc. Natl. Acad. Sci. USA* 1993; 90:6434-6438.
6. Hicks M, Delbridge L, Yue D, Reeve R. Catalysis of lipid peroxidation by glucose and glycosylated proteins. *Biochem. Biophys. Res. Commun.* 1988; 151:649-655.
7. Hunt J, Smith C, Wolff S. Autooxidative glycosylation and possible involvement of peroxides and free radicals in LDL modification by glucose. *Diabetes* 1990; 39:1420-1424.
8. Klein R, Klein B, Moss S, Davis M, DeMets DL. Glycosylated haemoglobin predicts the incidence and progression of diabetic retinopathy. *JAMA* 1989; 260:2864-2868.
9. Mullarkey C, Edelstein D, Brownlee M. Free radical generation by early glycation endproducts: a mechanism for accelerated atherogenesis in diabetes. *Biochem. Biophys. Res. Commun.* 1990; 173:932-939.
10. Sakurai T, Tsuchiya S. Superoxide production from nonenzymatically glycated protein. *FEBS Lett.* 1988; 236:406-410.
11. Schmidt AM, Vianna M, Gerlach M, et al. Isolation and characterization of binding proteins for advanced glycation endproducts from lung tissue which are present on the endothelial cell surface. *J. Biol. Chem.* 1992; 267:14987-14997.
12. Yang Z, Makita J, Hori Y, et al. Two novel rat liver membrane proteins that bind advanced glycosylation end products: relationship to macrophage receptor for glucose-modified proteins. *J. Exp. Med.* 1991; 174:515-524.
13. Neeper M, Schmidt AM, Brett J, Yan SD, Wang F, Pan YC, Elliston K, Stern D, Shaw A. Cloning and expression of RAGE: a cell surface receptor for advanced glycation endproducts of proteins. *J. Biol. Chem.* 1992; 267:14998-15004.
14. Nakayama H, Taneda S, Misuhashi T, Kuwajima S, Aoki S, Kuroda Y, Misawa K, Yanagisawa K, Nakagawa S. Characterization of antibodies to advanced glycosylation end products in vivo. *J. Immunol. Methods* 1991; 140:119-125.
15. Makita Z, Vlassara H, Cerami A, Bucala R. Immunochemical detection of advanced glycosylation end products in vivo. *J. Biol. Chem.* 1992; 267:5133-5138.
16. Schmidt AM, Yan SD, Brett J, Mora R, Nowygrod R, Stern D. Regulation of

hum:
adva:
17. Yan
Stern
glyc:
1994
18. Schr
1:10
19. Beis
link
beti
20. Dyc
lati:
Clir
21. Mc
lar:
dia:
22. Na
Ma
va:
dia:
23. Mi
N,
en
Cl
24. M
In
uc
vo
25. F
N
26. S
F
c
27. S
c
28. c
29.
30.

ular dis-
diabetes.
n human
iol Chem
oducts in
J. Med.
advanced
Sci. USA
lation by
un. 1988;
involve-
Diabetes
moglobin
4A 1989;
by early
diabetes.
glycated
ization of
ue which
17:14987-
teins that
ge recep-
, Stern D,
advanced
5004.
, Misawa
advanced
:119-125.
tection of
992; 267:
ulation of

- human mononuclear phagocyte migration by cell surface binding proteins for advanced glycation endproducts. *J. Clin. Invest.* 1993; 92:2155-2168.
17. Yan S-D, Schmidt A-M, Anderson G, Zhang J, Brett J, Zou Y-S, Pinsky D, Stern D. Enhanced cellular oxidant stress by the interaction of advanced glycation endproducts with their receptors/binding proteins. *J. Biol. Chem.* 1994; 269:9889-9897.
18. Schmidt AM, Yan SD, Stern D. The Dark Side of Glucose. *Nature Med.* 1995; 1:1002-1004.
19. Beisswenger P, Moore L, Brinck-Johnsen T, Curphey T. Increased collagen-linked pentosidine levels and advanced glycosylation endproducts in early diabetic nephropathy. *J. Clin. Invest.* 1991; 92:212-217.
20. Dyer D, Dunn J, Thorpe S, Bailie K, Lyons T, McCance D, Baynes J. Accumulation of Maillard reaction products in skin collagen in diabetes and aging. *J. Clin. Invest.* 1993; 91:2463-2469.
21. McCance D, Dyer D, Dunn D, Bailie K, Thorpe S, Baynes J, Lyons T. Maillard reaction products and their relation to complications in insulin-dependent diabetes mellitus. *J. Clin. Invest.* 1993; 91:2470-2478.
22. Nakamura Y, Horii Y, Nishino T, Shiki H, Sakaguchi Y, Kagoshima T, Dohi K, Makita Z, Vlassara H, Bucala R. Immunohistochemical localization of advanced glycosylation end products in coronary atheroma and cardiac tissue in diabetes mellitus. *Am. J. Pathol.* 1993; 143:1649-1656.
23. Miyata T, Oda O, Inagi R, Iida Y, Araki N, Yamada N, Horiuchi S, Taniguchi N, Maeda K, Kinoshita T. β_2 -Microglobulin modified with advanced glycation endproducts is a major component of hemodialysis-associated amyloidosis. *J. Clin. Invest.* 1993; 92:1243-1252.
24. Miyata T, Inagi R, Iida Y, Sato M, Yamada N, Oda O, Maeda K, Seo H. Involvement of β_2 -microglobulin modified with advanced glycation endproducts in the pathogenesis of hemodialysis-associated amyloidosis. *J. Clin. Invest.* 1994; 93:521-528.
25. Harrington C, Colaco CALS. Alzheimer's disease: a glycation connection. *Nature* 1994; 370:247-248.
26. Smith M, Kutty R, Richey P, Yan SD, Stern D, Chader G, Wiggert B, Petersen R, Perry G. Heme oxygenase-1 is associated with the neurofibrillary pathology of Alzheimer disease. *Am. J. Pathol.* 1994; 145:42-47.
27. Smith M, Taneda S, Richey P, Miyata S, Yan SD, Stern D, Monnier V, Perry G. Advanced Maillard reaction end products are associated with Alzheimer disease pathology. *Proc. Natl. Acad. Sci. USA* 1994; 91:5710-5714.
28. Yan SD, Chen X, Schmidt AM, et al. The presence of glycated tau in Alzheimer's disease: a mechanism for induction of oxidant stress. *Proc. Natl. Acad. Sci. USA* 1994; 91:7787-7791.
29. Vitek M, Bhattacharya K, Glendening JM. AGEs contribute to amyloidosis in Alzheimer's disease. *Proc. Natl. Acad. Sci. USA* 1994; 91:4766-4770.
30. Esposito C, Gerlach H, Brett J, Stern D, Vlassara H. Endothelial receptor-mediated binding of glucose-modified albumin is associated with increased monolayer permeability and modulation of cell surface coagulant properties. *J. Exp. Med.* 1989; 170:13887-1407.

31. Vlassara H, Bucala R, Striker L. Pathogenic effects of AGEs: Biochemical, biologic, and clinical implications for diabetes and aging. *Lab. Invest.* 1994; 70:138-151.
32. Khoury J, Thomas C, Loike J, Hickman S, Cao L, Silverstein S. Macrophages adhere to glucose-modified basement membranes via their scavenger receptors. *J. Biol. Chem* 1994; 269:10197-10200.
33. Hori O, Brett J, Slattery T, Cao R, Zhang J, Chen J, Nagashima M, Nitecki D, Morser J, Stern D, Schmidt A.M. The receptor for advanced glycation endproducts (RAGE) is a cellular binding site for amphoterin: mediation of neurite outgrowth and coexpression of RAGE and amphoterin in the developing nervous system. *J. Biol. Chem.* 1995; 270:25752-25761.
34. Brett J, Schmidt A-M, Zou Y-S, Yan S-D, Weidman E, Pinsky D, Neeper M, Przysiecki M, Shaw A, Migheli A, Stern D. Tissue distribution of the receptor for advanced glycation endproducts (RAGE): expression in smooth muscle, cardiac myocytes, and neural tissue in addition to the vasculature. *Am. J. Pathol.* 1993; 143:1699-1712.
35. Wautier JL, Wautier MP, Schmidt AM, Anderson GM, Hori O, Zoukourian C, Capron L, Chappey O, Brett J, Guillausseau PJ, Stern D. AGEs on the surface of diabetic erythrocytes bind to the vessel wall via a specific receptor inducing oxidant stress in the vasculature: a link between surface-associated AGEs and diabetic complications. *Proc Natl Acad Sci USA* 1994; 91:7742-7746.
36. Ross R. The pathogenesis of atherosclerosis—an update. *N. Engl. J. Med.* 1986; 314:488-500.
37. Kirstein M, Brett J, Radoff S, Ogawa S, Stern D, Vlassara H. Advanced protein glycosylation induces transendothelial human monocyte chemotaxis and secretion of platelet-derived growth factor: role in vascular disease of diabetes and aging. *Proc. Natl. Acad. Sci. USA* 1990; 87:9010-9014.
38. Vlassara H, Brownlee M, Manogue K, Dinarello C, Pasagian A. Cachectin/TNF and IL-1 induced by glucose-modified proteins: role in normal tissue remodeling. *Science* 1990; 240:1546-1548.
39. Schmidt AM, Hasu M, Popov D, et al. The receptor for advanced glycation endproducts (AGEs) has a central role in vessel wall interaction and gene activation in response to AGEs in the intravascular space. *Proc Natl Acad Sci USA* 1994; 91:8807-8811.
40. Schreck R, Rieber P, Baeuerle P. Reactive oxygen intermediates as apparently widely used messengers in the activation of the NF- κ B transcription factor and HIV-1. *EMBO J.* 1991; 10:2247-2258.
41. Collins T. Endothelial nuclear factor κ B and the initiation of the atherosclerotic lesion. *Lab Invest.* 1993; 10:336-349.
42. Marui, N, Offerman M, Swerlick R, et al. VCAM-1 gene transcription and expression are regulated through an oxidant-sensitive mechanism in human vascular endothelial cells. *J. Clin. Invest.* 1993; 92:1866-1984.
43. Cybulsky M, Gimbrone M. Endothelial expression of a mononuclear leukocyte adhesion molecule during atherogenesis. *Science* 1991; 251:788-791.
44. Li H, Cybulsky M, Gimbrone M, Libby P. An atherogenic diet rapidly induces

V
re
45. R
vi
a:
46. C
1
n
9.
47. S
E
e
(
to
48. N
t
49. I
c
1
50. S
v
c
c
51. P
c
t
52. 1
t
53.
54.
55.
56.

chemical,
est. 1994;

rophages
ger recep-

Nitecki D,
glycation
diation of
e develop-

Neeper M,
receptor
th muscle,
re. Am. J.

kourian C,
the surface
r inducing
AGEs and
6.
gl. J. Med.

Advanced
chemotaxis
disease of
4.
Cachectin/
rmal tissue

d glycation
1 and gene
tl Acad Sci

apparently
1 factor and

the athero-

ription and
1 in human

clear leuko-
8-791.
idly induces

VCAM-1: a cytokine-regulatable mononuclear leukocyte adhesion molecule in rabbit aortic endothelium. *Arterioscler. Thromb.* 1993; 13:197-204.

45. Richardson M, Hadcock S, DeReske, Cybulsky M. Increased expression in vivo of VCAM-1 and E-selectin by the aortic endothelium of normolipemic and hyperlipemic diabetic rabbits. *Arterioscler. Thromb.* 1994; 14:760-769.
46. O'Brien DK, Allen MD, McDonald TO, et al. Vascular cell adhesion molecule-1 is expressed in human coronary atherosclerotic plaques. Implications for the mode of progression of advanced coronary atherosclerosis. *J. Clin. Invest.* 1993; 92:945-951.
47. Schmidt A-M, Hori O, Chen J, Li J-F, Crandall J, Zhang J, Cao R, Yan S-D, Brett J, Stern D. Advanced glycation endproducts interacting with their endothelial receptor induce expression of vascular cell adhesion molecule-1 (VCAM-1): a potential mechanism for the accelerated vasculopathy of diabetes. *J. Clin. Invest.* 1995; 96:1395-1403.
48. Neish A, Williams H, Palmer H, Whitley M, Collins T. Functional analysis of the human VCAM-1 promoter. *J. Exp. Med.* 1992; 176:1583-1593.
49. Iademarco M, McQuillan J, Dean D. VCAM-1: contrasting transcriptional control mechanisms in muscle and endothelium. *Proc. Natl. Acad. Sci. USA* 1993; 90:3943-3947.
50. Schmidt AM, Crandall J, Cao R, Hori O, Lakatta E. Elevated plasma levels of vascular cell adhesion molecule-1 (VCAM-1) in diabetic patients with microalbuminuria: a marker of vascular dysfunction and progressive vascular disease. *Br. J. Hematol.* 1995; in press.
51. Messent J, Elliott T, Hill R, Jarrett R, Keen H, Viberti G. Prognostic significance of microalbuminuria in insulin-dependent diabetes mellitus: a twenty-three follow-up study. *Kidney Int* 1992; 41:836-839.
52. Nannipieri M, Rizzo L, Rapuano A, Pilo A, Penno G, Navelesi R. Increased transcapillary escape rate of albumin in microalbuminuric type II diabetic subject. *Diabetes Care* 1995; 18:1-9.
53. Koch AE, Halloran MM, Haskell CJ, Shah MR, Polverini PJ. Angiogenesis mediated by soluble forms of E-selectin and vascular cell adhesion molecule-1. *Nature* 1995; 376:517-519.
54. Wautier J-L, Zoukourian C, Chappey O, Wautier M-P, Guillausseau P-J, Cao R, Hori O, Stern D, Schmidt AM. Receptor-mediated endothelial cell dysfunction in diabetic vasculopathy: soluble receptor for advanced glycation endproducts blocks hyperpermeability. *J. Clin. Invest.* 1996; 97:238-243.
55. Williamson JR, Chang K, Tilton RG, Prater C, Jeffrey JR, Weigel C, Sherman WR. Increased vascular permeability in spontaneously diabetic BB/W rats and in rats with mild versus severe streptozocin-induced diabetes. *Diabetes* 1987; 36:813-821.
56. Albelda S, Sampson PM, Haselton FR, McNiff JM, Meller SN, Williams SK, Fishman AP. Permeability characteristics of cultured endothelial cell monolayers. *J. Appl. Physiol* 1988; 64:308-322.